

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books  
 Search PubMed for [inhibition[Title] AND tumor[Title] AND angiogenesis] Go Clear Advanced Search (beta) Save Search

Limits Preview/Index History Clipboard Details

Display  Show  Sort By  Send to   
All: 1 Review: 0

We found 1 articles:

1: Med Hypotheses. 2008;70(2):343-5. Epub 2007 Jul 13.

ELSEVIER  Links

### Inhibition of tumor angiogenesis by HMGB1 A box peptide.

Zhang CL, Shu MG, Qi HW, Li LW.

Department of Respiratory Medicine, Xijing Hospital, The Fourth Military Medical University, No. 17 Changle Western Road, Xi'an 710032, PR China.

High mobility group box 1 protein (HMG1) is a highly conserved, ubiquitous non-histone nuclear protein, which participates in maintaining nucleosome structure, regulation of gene transcription, and modulating the activity of steroid hormone receptors. Substantial evidence demonstrated that HMG1 could be secreted into the extracellular milieu, acts as a proinflammatory cytokine and mediates the chemotactic response of tumor cells to chemokines [1]. Recently, several reports suggested that HMG1 plays a key role in tumor angiogenesis through multiple mechanisms, including up-regulation of proangiogenic factors, promoting endothelial progenitor cells homing to ischemic tumor tissues and induction of endothelial cell migration and sprouting. And blockade of HMG1 binding to the receptor for advanced glycation end products (RAGE) could inhibit tumor angiogenesis [2-4]. RAGE, a transmembrane signaling antibody has been proved to inhibit angiogenesis efficiently. Since HMG1 A box peptide could antagonize the HMG1 whole length protein by competitively binding to RAGE and has been considered as a HMG1 specific antagonist, we postulate that the HMG1 A box peptide could function as an anti-angiogenic agent to inhibit tumor angiogenesis. In our opinion, if the hypothesis proved to be true, the HMG1 A box peptide could be widely used in clinical settings to treat malignant tumors.

PMID: 17630223 [PubMed - indexed for MEDLINE]

Display: AbstractPlus Show: 20 Sort By: Send to:

### Related Articles

### HMGB1 in sepsis

[Scand J Infect Dis, 2003]

**Convergence and amplification of toll-like receptor (TLR) and receptor for advanced glycation end products (RAGE) signaling pathways via high mobility group B1 (HMGB1).** (Angiogenesis: 2008)

**Cutting edge:** extracellular high mobility group box-1 protein is a proangiogenic cytokine [J Immunol. 2005]

High-mobility group box 1 activates integrin-dependent homing of endothelial progenitor cells. [Circ Res. 2007]

High mobility group 1 B-box mediates activation of human endothelium. [J Intern Med. 2003]

» [See all Related Articles.](#)

**Write to the Help Desk**

NCBI | NLM | NIH

Department of Health &amp; Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)